

## Dosing Information for Healthcare Professionals

Children's  
**MOTRIN**<sup>®</sup> Concentrated  
Infants' Drops

Use this chart to determine the proper dose of MOTRIN<sup>®</sup> for your patient. If possible, use weight to dose; otherwise use age.

**DOSE: Every 6-8 hours as needed.  
Do not give more than 4 doses in 24 hours.**

WEIGHT	AGE	Concentrated MOTRIN <sup>®</sup> Infants' Drops Active Ingredient: ibuprofen 50 mg (NSAID) <sup>*</sup> (in each 1.25 mL)	Children's MOTRIN <sup>®</sup> Oral Suspension Active Ingredient: ibuprofen 100 mg (NSAID) <sup>*</sup> (in each 5 mL or 1 tsp)
6-11 lbs	0-5 mos	—	—
12-17 lbs	6-11 mos	1.25 mL	—
18-23 lbs	12-23 mos	1.875 mL	—
24-35 lbs	2-3 yrs	—	5 mL (1 tsp)
36-47 lbs	4-5 yrs	—	7.5 mL (1½ tsp)
48-59 lbs	6-8 yrs	—	10 mL (2 tsp)
60-71 lbs	9-10 yrs	—	12.5 mL (2½ tsp)
72-95 lbs	11 yrs	—	15 mL (3 tsp)

mL = milliliter

tsp = teaspoon

### Remind parents and caregivers:

- Read and follow the label on all MOTRIN<sup>®</sup> products
- Take every 6-8 hours as needed
- Do NOT exceed more than 4 doses in 24 hours
- Use only the dosing device (syringe or dosing cup) that came with the product

<sup>\*</sup>Nonsteroidal anti-inflammatory drug

©McNEIL-PPC, Inc. 2014 • CTCM-0310