

Dosing Information for Healthcare Professionals

Infants'
TYLENOL *Children's*
TYLENOL

Use this chart to determine the proper dose of TYLENOL® for your patient. If possible, use weight to dose; otherwise use age.

WEIGHT	AGE	Infants' TYLENOL® Oral Suspension Active Ingredient: Acetaminophen 160 mg (in each 5 mL)	Children's TYLENOL® Oral Suspension Active Ingredient: Acetaminophen 160 mg (in each 5 mL or 1 tsp)
6-11 lbs	0-3 mos	1.25 mL	—
12-17 lbs	4-11 mos	2.5 mL	—
18-23 lbs	12-23 mos	3.75 mL	—
24-35 lbs	2-3 yrs	5 mL	5 mL (1 tsp)
36-47 lbs	4-5 yrs	—	7.5 mL (1½ tsp)
48-59 lbs	6-8 yrs	—	10 mL (2 tsp)
60-71 lbs	9-10 yrs	—	12.5 mL (2½ tsp)
72-95 lbs	11 yrs	—	15 mL (3 tsp)

mL = milliliter

tsp = teaspoon

**DOSE: Every 4 hours as needed.
Do not give more than 5 doses in 24 hours.**

Remind parents and caregivers:

- Read and follow the label on all TYLENOL® products
- Take every 4 hours as needed
- Do NOT exceed more than 5 doses in 24 hours
- Do NOT use with any other product containing acetaminophen
- Use only the dosing device (syringe or dosing cup) that came with the product

All Infants' TYLENOL® and Children's TYLENOL® Oral Suspension products have the same acetaminophen concentration (160 mg/5 mL)

©McNEIL-PPC, Inc. 2014 • CTCM-0310